

# AMS526: Numerical Analysis I (Numerical Linear Algebra for Computational and Data Sciences)

Lecture 10: Positive Definite Systems;  
Cholesky Factorization; Review for Midterm #1

Xiangmin Jiao

Stony Brook University

# Outline

1 Positive-Definite Systems (MC §4.2)

2 Cholesky Factorization (NL A§23)

3 Review of Midterm #1

# Symmetric Positive-Definite Matrices

- Symmetric matrix  $A \in \mathbb{R}^{n \times n}$  is *symmetric positive definite* (SPD) if  $x^T A x > 0$  for  $x \in \mathbb{R}^n \setminus \{0\}$
- Hermitian matrix  $A \in \mathbb{C}^{n \times n}$  is *Hermitian positive definite* (HPD) if  $x^* A x > 0$  for  $x \in \mathbb{C}^n \setminus \{0\}$
- SPD matrices have positive real eigenvalues and orthogonal eigenvectors
- Note: A positive-definite matrix does not need to be symmetric or Hermitian! A real matrix  $A$  is positive definite iff  $A + A^T$  is SPD
- If  $x^T A x \geq 0$  for  $x \in \mathbb{R}^n \setminus \{0\}$ , then  $A$  is said to be *positive semidefinite*

# Properties of Symmetric Positive-Definite Matrices

- SPD matrix often arises as Hessian matrix of some convex functional
  - ▶ E.g., least squares problems; partial differential equations
- If  $A$  is SPD, then  $A$  is nonsingular
- Let  $X$  be any  $n \times m$  matrix with full rank and  $n \geq m$ . Then
  - ▶  $X^T X$  is symmetric positive definite, and
  - ▶  $XX^T$  is symmetric positive semidefinite
- If  $A$  is  $n \times n$  SPD and  $X \in \mathbb{R}^{n \times m}$  has full rank and  $n \geq m$ , then  $X^T A X$  is SPD
- Any principal submatrix (picking some rows and corresponding columns) of  $A$  is SPD and  $a_{ii} > 0$

# Outline

- 1 Positive-Definite Systems (MC §4.2)
- 2 Cholesky Factorization (NL A§23)
- 3 Review of Midterm #1

# Cholesky Factorization

- If  $A$  is symmetric positive definite, then there is factorization of  $A$

$$A = R^T R,$$

where  $R$  is upper triangular, and all its diagonal entries are positive

- Key idea: take advantage and preserve symmetry and positive-definiteness during factorization
- Eliminate below diagonal and to the right of diagonal, we have

$$\begin{aligned} A &= \begin{bmatrix} a_{11} & b^T \\ b & K \end{bmatrix} = \begin{bmatrix} r_{11} & 0 \\ b/r_{11} & I \end{bmatrix} \begin{bmatrix} r_{11} & b^T/r_{11} \\ 0 & K - bb^T/a_{11} \end{bmatrix} \\ &= \begin{bmatrix} r_{11} & 0 \\ b/r_{11} & I \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & K - bb^T/a_{11} \end{bmatrix} \begin{bmatrix} r_{11} & b^T/r_{11} \\ 0 & I \end{bmatrix} = R_1^T A_1 R_1, \end{aligned}$$

where  $r_{11} = \sqrt{a_{11}}$ , where  $a_{11} > 0$

- $K - bb^T/a_{11}$  is principal submatrix of SPD  $A_1 = R_1^{-T} A R_1^{-1}$  and therefore is SPD, with positive diagonal entries

# Cholesky Factorization

- Apply recursively to obtain

$$A = \left( R_1^T R_2^T \cdots R_n^T \right) (R_n \cdots R_2 R_1) = R^T R, \quad r_{jj} > 0,$$

which is known as *Cholesky factorization*

- How to obtain  $R$  from  $R_n, \dots, R_2, R_1$ ? Recursively:

$$\begin{aligned} A &= \begin{bmatrix} r_{11} & 0 \\ s & I \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & A_1 \end{bmatrix} \begin{bmatrix} r_{11} & s^T \\ 0 & I \end{bmatrix} \\ &= \begin{bmatrix} r_{11} & 0 \\ s & I \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & \tilde{R}^T \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & \tilde{R} \end{bmatrix} \begin{bmatrix} r_{11} & s^T \\ 0 & I \end{bmatrix} \\ &= \begin{bmatrix} r_{11} & 0 \\ s & \tilde{R}^T \end{bmatrix} \begin{bmatrix} r_{11} & s^T \\ 0 & \tilde{R} \end{bmatrix} = R^T R \end{aligned}$$

- $R$  is “union” of  $k$ th rows of  $R_k$  ( $R^T$  is “union” of columns of  $R_k^T$ )
- Matrix  $A_1$  is called the *Schur complement* of  $a_{11}$  in  $A$

# Existence and Uniqueness

- Every SPD matrix has a unique Cholesky factorization
  - ▶ It exists because algorithm for Cholesky factorization always works for SPD matrices
  - ▶ Unique because once  $\alpha = \sqrt{a_{11}}$  is determined at each step, entire column  $w/\alpha$  is determined
- Question: How to check whether a symmetric matrix is positive definite?
- Answer: Run Cholesky factorization and it succeeds iff the matrix is positive definite.



# Algorithm of Cholesky Factorization

- Factorize SPD matrix  $A \in \mathbb{R}^{n \times n}$  into  $A = R^T R$

Algorithm: Cholesky factorization

$R = A$

**for**  $k = 1 : n$

**for**  $j = k + 1 : n$

$r_{j,j:n} \leftarrow r_{j,j:n} - (r_{kj}/r_{kk})r_{k,j:n}$

$r_{k,k:n} \leftarrow r_{k,k:n} / \sqrt{r_{kk}}$

- Note:  $r_{j,j:n}$  denotes subvector of  $j$ th row with columns  $j, j+1, \dots, n$
- Operation count

$$\sum_{k=1}^n \sum_{j=k+1}^n 2(n-j) \approx 2 \sum_{k=1}^n \sum_{j=1}^k j \approx \sum_{k=1}^n k^2 \approx \frac{n^3}{3}$$

- In practice,  $R$  overwrites  $A$ , and only upper-triangular part is stored.

# Notes on Cholesky Factorization

- Stability of Cholesky factorization
  - ▶ Cholesky factorization is backward stable
  - ▶ This is because  $\|R\|_2^2 = \|A\|_2$ , so entries in  $R$  are well bounded
- Cholesky factorization  $A = R^*R$  exists for HPD matrices, where  $R$  is upper-triangular and its diagonal entries are positive real values
- Implementations
  - ▶ Different versions of Cholesky factorization can all use block-matrix operators to achieve better performance, and actual performance depends on sizes of blocks
  - ▶ Different versions may have different amount of parallelism

## $LDL^T$ Factorization

- What happens if  $A$  is symmetric but not positive definite?
- Cholesky factorization is sometimes given by  $A = LDL^T$  where  $D$  is diagonal matrix and  $L$  is unit lower triangular matrix
- This avoids computing square roots
- Symmetric indefinite systems can be factorized with  $PAP^T = LDL^T$ , where
  - ▶  $P$  is a permutation matrix
  - ▶  $D$  is diagonal (if  $A$  is complex,  $D$  is block diagonal with  $1 \times 1$  and  $2 \times 2$  blocks)
  - ▶ its cost is similar to Cholesky factorization

# Banded Positive Definite Systems

- A matrix  $A$  is *banded* if there is a narrow band around the main diagonal such that all of the entries of  $A$  outside of the band are zero
- If  $A$  is  $n \times n$ , and there is an  $s \ll n$  such that  $a_{ij} = 0$  whenever  $|i - j| > s$ , then we say  $A$  is banded with bandwidth  $2s + 1$
- For symmetric matrices, only half of band is stored. We say that  $A$  has semi-bandwidth  $s$ .

## Theorem

*Let  $A$  be a banded, symmetric positive definite matrix with semi-bandwidth  $s$ . Then its Cholesky factor  $R$  also has semi-bandwidth  $s$ .*

- It is easy to prove using bordered form of Cholesky factorization
- Total flop count of Cholesky factorization is only  $\sim ns^2$
- However,  $A^{-1}$  of a banded matrix may be dense, so it is not economical to compute  $A^{-1}$

# Outline

- 1 Positive-Definite Systems (MC §4.2)
- 2 Cholesky Factorization (NL A§23)
- 3 Review of Midterm #1

# Midterm #1

- It will cover material up to Cholesky factorization
- It is a closed-book exam
- You can bring a single-sided, one-page, letter-size cheat sheet, which you must prepare by yourself

# Fundamental Concepts

- Norms, orthogonality, conditioning, stability
- Conditioning of problems
- Stability and backward stability of algorithms
- Efficiency of algorithms, operation counts
- Singular value decomposition, properties, and relationship with eigenvalue problems
- Orthogonal projection matrices, orthogonal matrices

# Algorithms

- Matrix multiplication
- Triangular systems
- Gaussian elimination with/without pivoting
- Cholesky factorization and  $LDL^T$  factorization
- Understand when they work, how they work, why they work, and how well they work